

(19)  Canadian
Intellectual Property
Office

An Agency of
Industry Canada

Office de la Propriété,
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

(11) **CA 2 479 766** (13) **A1**

(40) 02.10.2003
(43) 02.10.2003

(12)

(21) 2 479 766

(22) 19.03.2003

(51) Int. Cl. 7: **C07D 487/04, C07D 239/00,
C07D 249/00, C07C 255/57,
C07C 323/62, A01N 43/90**

(85) 17.09.2004

(86) PCT/EP03/002847

(87) WO03/080615

(30) 102 12 739.5 DE 21.03.2002
102 15 814.2 DE 10.04.2002
102 58 050.2 DE 11.12.2002

(71) BASF AKTIENGESELLSCHAFT,
67056, LUDWIGSHAFEN, XX (DE).

(72) LORENZ, GISELA (DE).
SCHOEFL, ULRICH (DE).
SCHWOEGLER, ANJA (DE).
MUELLER, BERND (DE).
BLETTNER, CARSTEN (DE).


SCHAEFER, PETER (DE).
GEWEHR, MARKUS (DE).
AMMERMANN, EBERHARD (DE).
GYPSER, ANDREAS (DE).
STRATHMANN, SIEGFRIED (DE).
GRAMMENOS, WASSILIOS (DE).
STIERL, REINHARD (DE).
GROTE, THOMAS (DE).
SCHIEWECK, FRANK (DE).
RHEINHEIMER, JOACHIM (DE).
TORMO I BLASCO, JORDI (DE).

(74) ROBIC

(54) TRIAZOLOPYRIMIDINES FONGICIDES, LEUR PROCEDE DE PRODUCTION ET LEUR UTILISATION POUR
LUTTER CONTRE DES CHAMPIGNONS NUISIBLES, ET AGENTS LES CONTENANT
(54) FUNGICIDAL TRIAZOLOPYRIMIDINES, METHODS FOR PRODUCING THE SAME, USE THEREOF FOR
COMBATING HARMFUL FUNGI AND AGENTS CONTAINING SAID SUBSTANCES

(57)

The invention relates to triazolopyrimidines of formula (I), in which the substituents are defined as follows: L1 represents cyano, S(=O)nA1 or C(=O)A2, wherein A1 stands for hydrogen, hydroxy, alkyl, alkylamino or dialkylamino; A2 stands for C1-C8 alkoxy, C1-C6 haloalkoxy or one of the groups named in A1; and n stands for 0, 1 or 2; L2, L3 represent hydrogen or halogen; L4, L5 represent hydrogen, halogen or alkyl; X represents halogen, cyano, alkyl, haloalkyl, alkoxy or haloalkoxy; R1 represents alkyl, haloalkyl, cycloalkyl, halocycloalkyl, alkenyl, alkadienyl, haloalkenyl, cycloalkenyl, alkynyl, haloalkynyl or cycloalkynyl, phenyl, naphthyl, or a five to ten-membered saturated, partially unsaturated or aromatic heterocyclous containing between one and four heteroatoms from the group containing O, N or S; R2 represents hydrogen or R1; R1 and R2 can form, together with the nitrogen atom to which they are bonded, a five or six-membered ring, which can be interrupted and/or substituted by an atom from the group O, N and S; whereby R1 and/or R2 can be substituted in accordance with the description. The invention also relates to methods and intermediate products for producing said compounds, to agents containing the latter and to the use of said compounds for combating harmful fungi.





Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2479766 A1 2003/10/02

(21) **2 479 766**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2003/03/19
(87) Date publication PCT/PCT Publication Date: 2003/10/02
(85) Entrée phase nationale/National Entry: 2004/09/17
(86) N° demande PCT/PCT Application No.: EP 2003/002847
(87) N° publication PCT/PCT Publication No.: 2003/080615
(30) Priorités/Priorities: 2002/03/21 (102 12 739.5) DE;
2002/04/10 (102 15 814.2) DE;
2002/12/11 (102 58 050.2) DE

(51) Cl.Int.⁷/Int.Cl.⁷ C07D 487/04, A01N 43/90, C07C 323/62,
C07D 239/00, C07C 255/57, C07D 249/00


(71) Demandeur/Applicant:
BASF AKTIENGESELLSCHAFT, DE

(72) Inventeurs/Inventors:
TORMO I BLASCO, JORDI, DE;
BLETTNER, CARSTEN, DE;
MUELLER, BERND, DE;
GEWEHR, MARKUS, DE;
GRAMMENOS, WASSILIOS, DE;
GROTE, THOMAS, DE;

...

(74) Agent: ROBIC

(54) Titre : TRIAZOLOPYRIMIDINES FONGICIDES, LEUR PROCEDE DE PRODUCTION ET LEUR UTILISATION POUR LUTTER CONTRE DES CHAMPIGNONS NUISIBLES, ET AGENTS LES CONTENANT
(54) Title: FUNGICIDAL TRIAZOLOPYRIMIDINES, METHODS FOR PRODUCING THE SAME, USE THEREOF FOR COMBATING HARMFUL FUNGI AND AGENTS CONTAINING SAID SUBSTANCES



(57) Abrégé/Abstract:

The invention relates to triazolopyrimidines of formula (I), in which the substituents are defined as follows: L¹ represents cyano, S(=O)_nA¹ or C(=O)A², wherein A¹ stands for hydrogen, hydroxy, alkyl, alkylamino or dialkylamino; A² stands for C₁-C₈ alkoxy, C₁-C₆ haloalkoxy or one of the groups named in A¹; and n stands for 0, 1 or 2; L², L³ represent hydrogen or halogen; L⁴, L⁵ represent hydrogen, halogen or alkyl; X represents halogen, cyano, alkyl, haloalkyl, alkoxy or haloalkoxy; R¹ represents alkyl, haloalkyl, cycloalkyl, halocycloalkyl, alkenyl, alkadienyl, haloalkenyl, cycloalkenyl, alkynyl, haloalkynyl or cycloalkynyl, phenyl, naphthyl, or a five to ten-membered saturated, partially unsaturated or aromatic heterocycclus containing between one and four heteroatoms from the group containing O, N or S; R² represents hydrogen or R¹; R¹ and R² can form, together with the nitrogen atom to which they are bonded, a five or six-membered ring, which can be interrupted and/or substituted by an atom from the group O, N and S; whereby R¹ and/or R² can be substituted in accordance with the description. The invention also relates to methods and intermediate products for producing said compounds, to agents containing the latter and to the use of said compounds for combating harmful fungi.

Canada

<http://opic.gc.ca> • Ottawa-Hull K1A 0C9 • <http://cipo.gc.ca>

OPIC • CIPO 191

OPIC



CIPO

(72) Inventeurs(suite)/Inventors(continued): GYPSE, ANDREAS, DE; RHEINHEIMER, JOACHIM, DE;
SCHAEFER, PETER, DE; SCHIEWECK, FRANK, DE; SCHWUEGLER, ANJA, DE; AMMERMAN, EBERHARD, DE;
STRATHMANN, SIEGFRIED, DE; LORENZ, GISELA, DE; STIERL, REINHARD, DE; SCHOEFL, ULRICH, DE

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG(19) Weltorganisation für geistiges Eigentum
Internationales Büro(43) Internationales Veröffentlichungsdatum
2. Oktober 2003 (02.10.2003)

PCT


(10) Internationale Veröffentlichungsnummer
WO 03/080615 A1

- (51) Internationale Patentklassifikation: C07D 487/04, A01N 43/90, C07C 255/57, 323/62 // (C07D 487/04, 249:00, 239:00)
- (71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme von US): BASF AKTIENGESELLSCHAFT [DE/DE]; 67056 Ludwigshafen (DE).
- (21) Internationales Aktenzeichen: PCT/EP03/02847
- (72) Erfinder; und (75) Erfinder/Anmelder (nur für US): TORMO I BLASCO, Jordi [DE/DE]; Carl-Benz-Str. 10-3, 69514 Laudenbach (DE). BLETNER, Carsten [DE/DE]; Richard-Wagner-Str. 48, 68165 Mannheim (DE). MÜLLER, Bernd [DE/DE]; Stockingerstr.7, 67227 Frankenthal (DE). GEWEHR, Markus [DE/DE]; Richard-Wagner-Strasse 2, 68165 Mannheim (DE). GRAMMENOS, Wasilios [DE/DE]; Samuel-Hahnemann Weg 9, 67071 Ludwigshafen (DE). GROTE, Thomas [DE/DE]; Im Hoehnhausen 18, 67157 Wachenheim (DE). GYPSE, Andreas [DE/DE]; B 4 4, 68159 Mannheim (DE). RHEINHEIMER, Joachim [DE/DE]; Merziger Str.24, 67063 Ludwigshafen (DE). SCHÄFER, Peter [DE/DE];
- (22) Internationales Anmeldedatum: 19. März 2003 (19.03.2003)
- (25) Einreichungssprache: Deutsch
- (26) Veröffentlichungssprache: Deutsch
- (30) Angaben zur Priorität:
102 12 739.5 21. März 2002 (21.03.2002) DE
102 15 814.2 10. April 2002 (10.04.2002) DE
102 58 050.2 11. Dezember 2002 (11.12.2002) DE

[Fortsetzung auf der nächsten Seite]

(54) Title: FUNGICIDAL TRIAZOLOPYRIMIDINES, METHODS FOR PRODUCING THE SAME, USE THEREOF FOR COMBATING HARMFUL FUNGI AND AGENTS CONTAINING SAID SUBSTANCES

(54) Bezeichnung: FUNGIZIDE TRIAZOLOPYRIMIDINE, VERFAHREN ZU IHRER HERSTELLUNG UND IHRE VERWENDUNG ZUR BEKÄMPFUNG VON SCHADPILZEN SOWIE SIE ENTHALTENDE MITTEL



(57) Abstract: The invention relates to triazolopyrimidines of formula (I), in which the substituents are defined as follows: L¹ represents cyano, S(=O)_nA¹ or C(=O)A², wherein A¹ stands for hydrogen, hydroxy, alkyl, alkylamino or dialkylamino; A² stands for C₁-C₈ alkoxy, C₁-C₆ haloalkoxy or one of the groups named in A¹; and n stands for 0, 1 or 2; L², L³ represent hydrogen or halogen; L⁴, L⁵ represent hydrogen, halogen or alkyl; X represents halogen, cyano, alkyl, haloalkyl, alkoxy or haloalkoxy; R¹ represents alkyl, haloalkyl, cycloalkyl, halocycloalkyl, alkenyl, alkadienyl, haloalkenyl, cycloalkenyl, alkynyl, haloalkynyl or cycloalkynyl, phenyl, naphthyl, or a five to ten-membered saturated, partially unsaturated or aromatic heterocyclo containing between one and four heteroatoms from the group containing O, N or S; R² represents hydrogen or R¹; R¹ and R² can form, together with the nitrogen atom to which they are bonded, a five or six-membered ring, which can be interrupted and/or substituted by an atom from the group O, N and S; whereby R¹ and/or R² can be substituted in accordance with the description. The invention also relates to methods and intermediate products for producing said compounds, to agents containing the latter and to the use of said compounds for combating harmful fungi.

(57) Zusammenfassung: Triazolopyrimidine der Formel (I), in der die Substituenten folgende Bedeutung haben: L¹ Cyano, S(=O)_nA¹ oder C(=O)A², worin A¹ Wasserstoff, Hydroxy, Alkyl, Alkylamino oder Di-alkylamino; A² C₁-C₈-Alkoxy, C₁-C₆-Halogenalkoxy oder eine der bei A¹ genannten Gruppen; n 0, 1 oder 2; L², L³ Wasserstoff oder Halogen; L⁴, L⁵ Wasserstoff, Halogen oder Alkyl; X Halogen, Cyano, Alkyl, Halogenalkyl, Alkoxy oder Halogenalkoxy; R¹ Alkyl, Halogenalkyl, Cycloalkyl, Halogenalkyl, Alkenyl, Alkadienyl, Halogenalkenyl, Cycloalkenyl, Alkynyl, Halogenalkynyl oder Cycloalkynyl, Phenyl, Naphthyl, oder ein fünf- bis zehngliedriger gesättigter, partiell ungesättigter oder aromatischer Heterocyclus, enthaltend ein bis vier Heteroatome aus der Gruppe O, N oder S; R² Wasserstoff oder R¹; R¹ und R² können auch zusammen mit dem Stickstoffatom, an das sie gebunden sind, einen fünf- oder sechsgliedrigen Ring bilden, der durch ein Atom aus der Gruppe O, N und S unterbrochen sein und/oder substituiert sein; wobei R¹ und/oder R² gemäß der Beschreibung substituiert sein können; Verfahren und Zwischenprodukte zur Herstellung dieser Verbindungen, sie enthaltende Mittel sowie ihre Verwendung zur Bekämpfung von Schadpilzen.

WO 03/080615 A1

WO 03/080615 A1



Römerstr.1, 67308 Ottersheim (DE). **SCHIEWECK, Frank** [DE/DE]; Lindenweg 4, 67258 Hessheim (DE). **SCHWÖGLER, Anja** [DE/DE]; Heinrich-Lanz-Str. 3, 68165 Mannheim (DE). **AMMERMAN, Eberhard** [DE/DE]; Von-Gagem-Str.2, 64646 Heppenheim (DE). **STRATHMANN, Siegfried** [DE/DE]; Donnersbergstr.9, 67117 Limburgerhof (DE). **LORENZ, Gisela** [DE/DE]; Erlenweg 13, 67434 Neustadt (DE). **STIERL, Reinhard** [DE/DE]; Jahnstr. 8, 67251 Freinsheim (DE). **SCHÖFL, Ulrich** [DE/DE]; Luftschiffing 22c, 68782 Brühl (DE).

MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(74) **Gemeinsamer Vertreter:** **BASF AKTIENGESSELLSCHAFT**; 67056 Ludwigshafen (DE).

(81) **Bestimmungsstaaten (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,

(84) **Bestimmungsstaaten (regional):** ARIPO-Patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI-Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Veröffentlicht:


— mit internationalem Recherchenbericht

Zur Erklärung der Zweibuchstaben-Codes und der anderen Abkürzungen wird auf die Erklärungen ("Guidance Notes on Codes and Abbreviations") am Anfang jeder regulären Ausgabe der PCT-Gazette verwiesen.

**FUNGICIDAL TRIAZOLOPYRIMIDINES, METHODS FOR PRODUCING THE
SAME, USE THEREOF FOR COMBATING HARMFUL FUNGI AND AGENTS
CONTAINING SAID SUBSTANCES**

5 The present invention relates to triazolopyrimidines of the formula I,

10




I

in which substituents have the following meanings:

- 15 L^1 is cyano, $S(=O)_nA^1$ or $C(=O)A^2$, wherein
- A^1 is hydrogen, hydroxyl, C_1 - C_8 -alkyl,
 C_1 - C_8 -alkylamino or di(C_1 - C_8 -alkyl)amino;
- 20 A^2 is C_1 - C_8 -alkoxy, C_1 - C_6 -haloalkoxy or one of the
 groups mentioned under A^1 ;
- n is 0, 1 or 2;
- 25 L^2, L^3 are hydrogen or halogen;
- L^4, L^5 are hydrogen, halogen or C_1 - C_4 -alkyl;
- 30 X is halogen, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl,
 C_1 - C_4 -alkoxy or C_1 - C_2 -haloalkoxy;
- R^1 is C_1 - C_8 -alkyl, C_1 - C_8 -haloalkyl, C_3 - C_6 -cycloalkyl,
 C_3 - C_6 -halocycloalkyl, C_2 - C_8 -alkenyl, C_4 - C_{10} -alkadienyl,
 C_2 - C_8 -haloalkenyl, C_3 - C_6 -cycloalkenyl, C_2 - C_8 -alkynyl,
35 C_2 - C_8 -haloalkynyl or C_3 - C_6 -cycloalkynyl, phenyl,
 naphthyl, or a five- to ten-membered saturated,
 partially unsaturated or aromatic heterocycle,
 comprising one to four heteroatoms from the group
 consisting of O, N and S;
- 40 R^2 is hydrogen or one of the groups mentioned under R^1 ,

1a

R¹ and R² can also, together with the nitrogen atom to which they are bonded, form a five- or six-membered ring which can be interrupted by an atom from the group consisting of O, N and S and/or can carry one or more substituents from the group consisting of halogen,



2

C₁-C₆-alkyl, C₁-C₆-haloalkyl and oxy-C₁-C₃-alkylenoxy or in which an N and a neighboring C atom can be connected via a C₁-C₄-alkylene chain;

5 wherein R¹ and/or R² can be substituted by one to four identical or different groups R^a:

10 R^a is halogen, cyano, nitro, hydroxyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkylcarbonyl, C₃-C₆-cycloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₁-C₆-alkoxycarbonyl, C₁-C₆-alkylthio, C₁-C₆-alkylamino, di(C₁-C₆-alkyl)amino, C₂-C₆-alkenyl, C₂-C₆-alkenyloxy, C₃-C₆-alkynyloxy, C₃-C₆-cycloalkyl, phenyl, naphthyl or a five- to
15 ten-membered saturated, partially unsaturated or aromatic heterocycle, comprising one to four heteroatoms from the group consisting of O, N and S,

20 wherein these aliphatic, alicyclic or aromatic groups, for their part, can be partially or completely halogenated or can carry one to three groups R^b:

25 R^b is halogen, cyano, nitro, hydroxyl, mercapto, amino, carboxyl, aminocarbonyl, aminothiocabonyl, alkyl, haloalkyl, alkenyl, alkenyloxy, alkynyloxy, alkoxy, haloalkoxy, alkylthio, alkylamino, dialkylamino, formyl, alkylcarbonyl, alkylsulfonyl, alkylsulfoxyl, alkoxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminothiocabonyl or
30 dialkylaminothiocabonyl, wherein the alkyl groups in these radicals comprise 1 to 6 carbon atoms and the alkenyl or alkynyl groups mentioned in these radicals comprise 2 to
35 8 carbon atoms;

40 and/or one to three of the following radicals:

cycloalkyl, cycloalkoxy, heterocyclyl or heterocyclxyloxy, wherein the cyclic systems
comprise 3 to 10 ring members; aryl, aryloxy, arylthio, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, hetaryl, hetaryloxy or hetarylthio, wherein the
45 aryl radicals preferably comprise 6 to 10 ring

3

members and the hetaryl radicals comprise 5 or 6 ring members, wherein the cyclic systems can be partially or completely halogenated or can be substituted by alkyl or haloalkyl groups.

5

The invention relates in addition to processes for and intermediates in the preparation of these compounds, compositions comprising them and their use in the control of harmful fungi.

10 5-Chlorotriazolopyrimidines for the control of harmful fungi are known from EP-A 71 792, EP-A 550 113, WO-A 98/46608 and WO-A 99/41255.

Fungicidally effective triazolopyrimidines having specific
15 substitution of the 6-phenyl group are known from EP-A 834 513, WO 98/46607, EP-A 945 453, WO 99/48893, US 5 985 883 and WO 02/46195.

However, in many cases, in particular when low amounts are
20 applied, their action is not always satisfactory.

It is the object of the present invention to provide compounds with an improved action and/or a broadened spectrum of activity.

25 We have found that this object is achieved with the compounds defined at the start. Processes for and intermediates in their preparation, compositions comprising them and processes for the control of harmful fungi through the use of the compounds I have also been found.

30


The compounds of the formula I are distinguished from those from the abovementioned documents in the substitution of the 6-phenyl group, which carries a group L¹ in the para-position, and the substitution of the 7-amino group.

35

The compounds of the formula I have, in comparison with the known compounds, an increased activity against harmful fungi.

The compounds according to the invention can be obtained in
40 various ways. They are advantageously prepared by reaction of 5-aminotriazole of the formula II with appropriately substituted phenylmalonates of the formula III in which R is alkyl, preferably C₁-C₆-alkyl, in particular methyl or ethyl.

45



This reaction is usually carried out at temperatures of 80°C to 250°C, preferably 120°C to 180°C, without solvent or in an inert organic solvent, in the presence of a base [cf. EP-A 770 615] or in the presence of acetic acid under the conditions known from Adv. Het. Chem., Vol. 57, pp. 81ff. (1993).

Suitable solvents are aliphatic hydrocarbons, aromatic hydrocarbons, such as toluene or o-, m- and p-xylene, halogenated hydrocarbons, ethers, nitriles, ketones, alcohols, N-methylpyrrolidone, dimethyl sulfoxide, dimethylformamide and dimethylacetamide. In a particularly preferred way, the reaction is carried out without solvent or in chlorobenzene, xylene, dimethyl sulfoxide, N-methylpyrrolidone. Mixtures of the solvents mentioned can also be used.

Suitable bases are generally inorganic compounds, such as alkali metal and alkaline earth metal hydroxides, alkali metal and alkaline earth metal oxides, alkali metal and alkaline earth metal hydrides, alkali metal amides, alkali metal and alkaline earth metal carbonates, alkali metal hydrogencarbonates, or organometallic compounds, in particular alkali metal alkyls, alkyl magnesium halides, and alkali metal and alkaline earth metal alkoxides and magnesium dimethoxide, as well as organic bases, e.g. tertiary amines, such as trimethylamine, triethylamine, triisopropylethylamine, tributylamine, N-methylpiperidine and N-methylmorpholine, pyridine, substituted pyridines, such as collidine, lutidine and 4-dimethylaminopyridine, and bicyclic amines. Tertiary amines, such as triisopropylethylamine, tributylamine, N-methylmorpholine or N-methylpiperidine, are especially preferred.

The bases are generally used in catalytic amounts. However, they can also be used in equimolar amounts, in excess or possibly as solvent.


The starting materials are generally reacted with one another in equimolar amounts. It can be advantageous to the yield to use the base and the malonate III in an excess with respect to the triazole.

5

Phenylmalonates of the formula III are advantageously obtained from the reaction of suitably substituted bromobenzenes with dialkyl malonates under Cu(I) catalysis [cf. Chemistry Letters, pp. 367-370, 1981; EP-A 10 02 788].

5


The dihydroxytriazolopyrimidines of the formula IV are converted to the dihalopyrimidines of the formula V under the conditions known from WO-A 94/20501. A chlorinating agent or a brominating agent, such as phosphorus oxybromide or phosphorus oxychloride, optionally in the presence of a solvent, is advantageously used as halogenating agent [Hal].



This reaction is usually carried out at 0°C to 150°C, preferably at 80°C to 125°C [cf. EP-A 770 615].

20

Dihalopyrimidines of the formula V are further reacted with amines of the formula VI,



in which R¹ and R² are defined as in formula I, to give compounds of the formula I in which X is halogen.

30


This reaction is advantageously carried out at 0°C to 70°C, preferably 10°C to 35°C, preferably in the presence of an inert solvent, such as ethers, e.g. dioxane, diethyl ether or, particularly, tetrahydrofuran, halogenated hydrocarbons, such as dichloromethane, and aromatic hydrocarbons, such as, for example, toluene [cf. WO-A 98/46608].

The use of a base, such as tertiary amines, for example triethylamine, or inorganic amines, such as potassium carbonate, is preferred; in addition, excess amine of the formula VI can be used as a base.

Compounds of the formula I in which X represents cyano, C₁-C₆-alkoxy or C₁-C₂-haloalkoxy can advantageously be obtained from the reaction of compounds I, in which X represents halogen, preferably chlorine, with compounds M-X' (formula VII). Compounds VII represent, depending on the meaning of the group X' to be

6


introduced, an inorganic cyanide, an alkoxide or a haloalkoxide. The reaction is advantageously carried out in the presence of an inert solvent. The cation M in formula VII is of little importance; for practical reasons, ammonium, tetraalkylammonium, 5 alkali metal or alkaline earth metal salts are usually preferred.



10 The reaction temperature usually lies between 0 and 120°C, preferably between 10 and 40°C [cf. J. Heterocycl. Chem., Vol.12, pp. 861-863 (1975)].

Suitable solvents include ethers, such as dioxane, diethyl ether 15 and, preferably, tetrahydrofuran, halogenated hydrocarbons, such as dichloromethane, and aromatic hydrocarbons, such as toluene.

Compounds of the formula I in which X is C₁-C₄-alkyl or C₁-C₄-haloalkyl can advantageously be obtained through the 20 following synthetic route:



The 5-alkyl-7-hydroxy-6-phenyltriazaolopyrimidines IVa are obtained from the ketoesters IIIa. The 5-methyl-7-hydroxy- 30 6-phenyltriazaolopyrimidines are obtained through the use of the readily accessible 2-phenylacetoacetic esters (IIIa with X¹=CH₃) [cf. Chem. Pharm. Bull., 9, 801, (1961)]. The preparation of the starting compounds IIIa is advantageously carried out under the conditions described in EP-A 10 02 788.

35

The 5-alkyl-7-hydroxy-6-phenyltriazaolopyrimidines obtained in this way are reacted with halogenating agents [Hal] to give the 7-halotriazaolopyrimidines of the formula Va. Chlorinating or brominating agents, such as phosphorus oxybromide, phosphorus 40 oxychloride, thionyl chloride, thionyl bromide or sulfur chloride, are preferably used. The reaction can be carried out in bulk or in the presence of a solvent. Usual reaction temperatures are from 0 to 150°C or, preferably, from 80 to 125°C.

45

7



10

15 C₁-C₃-alkyl and R represents C₁-C₄-alkyl. They are reacted to give compounds of the formula IX and are decarboxylated to give compounds I [cf. US 5,994,360].



30

35


40

45

8

pentane, hexane, cyclohexane and petroleum ether, aromatic hydrocarbons, such as toluene and o-, m- and p-xylene, halogenated hydrocarbons, such as methylene chloride, chloroform and chlorobenzene, ethers, such as diethyl ether, diisopropyl ether, tert-butyl methyl ether, dioxane, anisole and tetrahydrofuran, nitriles, such as acetonitrile and propionitrile, ketones, such as acetone, methyl ethyl ketone, diethyl ketone and tert-butyl methyl ketone, alcohols, such as methanol, ethanol, n-propanol, isopropanol, n-butanol and tert-butanol, dimethyl sulfoxide, dimethylformamide and dimethylacetamide. In a particularly preferred way, the reaction is carried out in hydrochloric acid or acetic acid. Mixtures of the solvents mentioned can also be used.

- 15 Compounds of the formula I in which X is C₁-C₄-alkyl can also be obtained by coupling of 5-halotriazolopyrimidines of the formula I in which X represents halogen with organometallic reagents of the formula X. In an embodiment of this process, the reaction is carried out under transition metal catalysis, such as Ni or Pd
- 20 catalysis.



- 25 In formula X, M is a metal ion with a valency of X, such as, for example, B, Zn or Sn. This reaction can be carried out, for example, analogously to the following methods: J. Chem. Soc. Perkin Trans., 1, 1187 (1994), ibid, 1, 2345 (1996); WO-A 99/41255; Aust. J. Chem., Vol. 43, 733 (1990); J. Org. Chem., Vol. 43, 358 (1978); J. Chem. Soc. Chem. Commun., 866 (1979); Tetrahedron Lett., Vol. 34, 8267 (1993); ibid, Vol. 33, 413 (1992).

If R¹ and R² represent halogen-free groups, optically active amines of the formula VI in the (R)-configuration are preferred.

If R¹ or R² comprises haloalkyl or haloalkenyl groups, the (S)-configuration is preferred for optically active amines of the formula VI.

- 40 Amines of the formula VI are either commercially available or, if they have a center of chirality, are accessible through resolution according to WO 02/38565. For example, (R)-3,3-dimethylbut-2-amine (R-DMBA) can particularly advantageously be prepared in this way.
- 45

9

In a preferred embodiment of the process according to the invention for the preparation of the compounds in which L^1 represents $S(=O)_nA^1$ with $n = 1$ or 2 , the thio compounds with $n = 0$ are oxidized at the formula I stage [Lit.: WO 94/14761; Synth, 5 Commun., Vol. 16, p. 233 (1986)].

Compounds of the formulae I, III and IV in which L^1 is $S(=O)_nA^1$ with $n = 2$ and A^1 is alkyl, in particular methyl, can also be used as intermediates in the preparation of additional
10 6-phenyltriazolopyrimidines. These intermediates can advantageously be used for the preparation of those triazolopyrimidines in which the 6-phenyl group exhibits in the para-position a group which can be introduced as a nucleophile, such as, for example, cyano, nitro, hydroxyl, alkoxy, haloalkoxy
15 or groups bonded via nitrogen, such as alkylamino, dialkylamino or a heterocycle bonded via N [Lit.: Tetrahedron Lett., p. 759 (1967); ibid p. 1763 (2000); J. Org. Chem., p. 4705 (1979)]. The exchange of the SO_2 alkyl group with the substituent to be introduced takes place particularly advantageously at the formula
20 I stage.

Compounds of the formula I in which L^1 is $C(=O)A^2$ with $A^2 =$ hydrogen or alkyl are advantageously prepared from the corresponding compounds in which L^1 is CN. This conversion is
25 carried out particularly preferably at the formula I stage.

Compounds in which L^1 represents CHO are preferably prepared from the corresponding cyanides by reduction under known conditions [cf. Collect. Czech. Chem. Commun., p. 729 (2000); J. Org. Chem.,
30 p. 5298 (2000); Heterocycles, p. 1173 (1987); Chem. Pharm. Bull., p. 1440 (1991)]. Compounds in which L^1 represents $C(O)alkyl$ are advantageously obtained from the corresponding cyanides by reaction with Grignard or alkyllithium compounds under known conditions [cf. J. Org. Chem., p. 4844 (1994); Synthetic Commun.,
35 p. 4067 (1998); Tetrahedron Lett., p. 6505 (1988)].

The reaction mixtures are worked up conventionally, e.g. by mixing with water, separating the phases and possibly chromatographic purification of the crude products. Some of the
40 intermediates and final products are obtained in the form of colorless or slightly brownish viscous oils which, under reduced pressure and at moderately elevated temperature, are freed or purified from volatile constituents. If the intermediates and final products are obtained as solids, the purification can also
45 take place by recrystallization or trituration.

10

If individual compounds I are not accessible by the methods described above, they can be prepared by derivatization of other compounds I.

- 5 If isomeric mixtures are obtained in the synthesis, a separation is, however, generally not absolutely necessary, since the individual isomers can sometimes be converted into one another during preparation for application or upon application (e.g., under the effect of light, acid or bases). Corresponding
10 conversions can also take place after application, for example, in the treatment of plants, in the treated plant or in the harmful fungus to be controlled.

Collective terms were used in the definitions of the symbols
15 given in the above formulae, which collective terms are generally representative of the following substituents:

halogen: fluorine, chlorine, bromine and iodine;

- 20 **alkyl:** saturated, straight-chain or branched hydrocarbon radicals with 1 to 4, 6 or 8 carbon atoms, e.g. C₁-C₆-alkyl, such as methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl,
25 hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl,
30 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl;

- haloalkyl:** straight-chain or branched alkyl groups with 1 to 8 carbon atoms (as mentioned above), in which the hydrogen atoms in
35 these groups can be partially or completely replaced by halogen atoms as mentioned above, e.g. C₁-C₂-haloalkyl, such as chloromethyl, bromomethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl,
40 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl, pentafluoroethyl or
1,1,1-trifluoroprop-2-yl;

45

11

- alkenyl:** unsaturated, straight-chain or branched hydrocarbon radicals with 2 to 4, 6 or 8 carbon atoms and a double bond in any position, e.g. C₂-C₆-alkenyl, such as ethenyl, 1-propenyl, 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 5 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-1-butenyl, 2-methyl-1-butenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 10 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-1-propenyl, 1-ethyl-2-propenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-1-pentenyl, 2-methyl-1-pentenyl, 3-methyl-1-pentenyl, 4-methyl-1-pentenyl, 15 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 20 1,2-dimethyl-1-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-1-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-1-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 25 3,3-dimethyl-1-butenyl, 3,3-dimethyl-2-butenyl, 1-ethyl-1-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-1-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl, 1-ethyl-2-methyl-1-propenyl and 1-ethyl-2-methyl-2-propenyl; 30
- haloalkenyl:** unsaturated, straight-chain or branched hydrocarbon radicals with 2 to 8 carbon atoms and a double bond in any position (as mentioned above), in which the hydrogen atoms in these groups can be partially or completely replaced by halogen 35 atoms as mentioned above, in particular fluorine, chlorine and bromine;

- alkynyl:** straight-chain or branched hydrocarbon groups with 2 to 4, 6 or 8 carbon atoms and a triple bond in any position, e.g. 40 C₂-C₆-alkynyl, such as ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 1-methyl-2-propynyl, 1-pentyne, 2-pentyne, 3-pentyne, 4-pentyne, 1-methyl-2-butyne, 1-methyl-3-butyne, 2-methyl-3-butyne, 3-methyl-1-butyne, 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 1-hexynyl, 45 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 1-methyl-2-pentyne, 1-methyl-3-pentyne, 1-methyl-4-pentyne, 2-methyl-3-pentyne, 2-methyl-4-pentyne, 3-methyl-1-pentyne, 3-methyl-4-pentyne,

12

4-methyl-1-pentynyl, 4-methyl-2-pentynyl, 1,1-dimethyl-2-butynyl,
1,1-dimethyl-3-butynyl, 1,2-dimethyl-3-butynyl,
2,2-dimethyl-3-butynyl, 3,3-dimethyl-1-butynyl,
1-ethyl-2-butynyl, 1-ethyl-3-butynyl, 2-ethyl-3-butynyl and
5 1-ethyl-1-methyl-2-propynyl;

haloalkynyl: unsaturated, straight-chain or branched hydrocarbon
radicals with 2 to 8 carbon atoms and a triple bond in any
position (as mentioned above), in which the hydrogen atoms in
10 these groups can be partially or completely replaced by halogen
atoms as mentioned above, in particular fluorine, chlorine and
bromine;

cycloalkyl: saturated mono- or bicyclic hydrocarbon groups with 3
15 to 6 or 8 carbon ring members, e.g. C₃-C₈-cycloalkyl, such as
cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and
cyclooctyl;

alkoxycarbonyl: an alkoxy group with 1 to 6 carbon atoms (as
20 mentioned above) which is bonded to the backbone via a carbonyl
group (-CO-);

oxyalkylenoxy: unbranched divalent chains formed from 1 to 3 CH₂
groups in which both valencies is bonded to the backbone via an
25 oxygen atom, e.g. OCH₂O, OCH₂CH₂O and OCH₂CH₂CH₂O;

**five- to ten-membered saturated, partially unsaturated or
aromatic heterocycle comprising one to four heteroatoms from the
group consisting of O, N and S:**
30 - 5- or 6-membered heterocyclyl comprising one to three nitrogen
atoms and/or one oxygen or sulfur atom or one or two oxygen
and/or sulfur atoms, e.g. 2-tetrahydrofuranyl,
3-tetrahydrofuranyl, 2-tetrahydrothienyl, 3-tetrahydrothienyl,
2-pyrrolidinyl, 3-pyrrolidinyl, 3-isoxazolidinyl,
35 4-isoxazolidinyl, 5-isoxazolidinyl, 3-isothiazolidinyl,
4-isothiazolidinyl, 5-isothiazolidinyl, 3-pyrazolidinyl,
4-pyrazolidinyl, 5-pyrazolidinyl, 2-oxazolidinyl,
4-oxazolidinyl, 5-oxazolidinyl, 2-thiazolidinyl,
4-thiazolidinyl, 5-thiazolidinyl, 2-imidazolidinyl,
40 4-imidazolidinyl, 1,2,4-oxadiazolidin-3-yl,
1,2,4-oxadiazolidin-5-yl, 1,2,4-thiadiazolidin-3-yl,
1,2,4-thiadiazolidin-5-yl, 1,2,4-triazolidin-3-yl,
1,3,4-oxadiazolidin-2-yl, 1,3,4-thiadiazolidin-2-yl,
1,3,4-triazolidin-2-yl, 2,3-dihydrofur-2-yl,
45 2,3-dihydrofur-3-yl, 2,4-dihydrofur-2-yl, 2,4-dihydrofur-3-yl,
2,3-dihydrothien-2-yl, 2,3-dihydrothien-3-yl,
2,4-dihydrothien-2-yl, 2,4-dihydrothien-3-yl, 2-pyrrolin-2-yl,

13

- 2-pyrrolin-3-yl, 3-pyrrolin-2-yl, 3-pyrrolin-3-yl,
2-isoxazolin-3-yl, 3-isoxazolin-3-yl, 4-isoxazolin-3-yl,
2-isoxazolin-4-yl, 3-isoxazolin-4-yl, 4-isoxazolin-4-yl,
2-isoxazolin-5-yl, 3-isoxazolin-5-yl, 4-isoxazolin-5-yl,
5 2-isothiazolin-3-yl, 3-isothiazolin-3-yl, 4-isothiazolin-3-yl,
2-isothiazolin-4-yl, 3-isothiazolin-4-yl, 4-isothiazolin-4-yl,
2-isothiazolin-5-yl, 3-isothiazolin-5-yl, 4-isothiazolin-5-yl,
2,3-dihydropyrazol-1-yl, 2,3-dihydropyrazol-2-yl,
2,3-dihydropyrazol-3-yl, 2,3-dihydropyrazol-4-yl,
10 2,3-dihydropyrazol-5-yl, 3,4-dihydropyrazol-1-yl,
3,4-dihydropyrazol-3-yl, 3,4-dihydropyrazol-4-yl,
3,4-dihydropyrazol-5-yl, 4,5-dihydropyrazol-1-yl,
4,5-dihydropyrazol-3-yl, 4,5-dihydropyrazol-4-yl,
4,5-dihydropyrazol-5-yl, 2,3-dihydrooxazol-2-yl,
15 2,3-dihydrooxazol-3-yl, 2,3-dihydrooxazol-4-yl,
2,3-dihydrooxazol-5-yl, 3,4-dihydrooxazol-2-yl,
3,4-dihydrooxazol-3-yl, 3,4-dihydrooxazol-4-yl,
3,4-dihydrooxazol-5-yl, 3,4-dihydrooxazol-2-yl,
3,4-dihydrooxazol-3-yl, 3,4-dihydrooxazol-4-yl, 2-piperidinyl,
20 3-piperidinyl, 4-piperidinyl, 1,3-dioxan-5-yl,
2-tetrahydropyranyl, 4-tetrahydropyranyl, 2-tetrahydrothienyl,
3-hexahydropyridazinyl, 4-hexahydropyridazinyl,
2-hexahydropyrimidinyl, 4-hexahydropyrimidinyl,
5-hexahydropyrimidinyl, 2-piperazinyl,
25 1,3,5-hexahydrotriazin-2-yl and 1,2,4-hexahydrotriazin-3-yl;
- 5-membered heteroaryl comprising one to four nitrogen atoms or
one to three nitrogen atoms and one sulfur or oxygen atom:
5-ring heteroaryl groups which, in addition to carbon atoms,
can comprise one to four nitrogen atoms or one to three
30 nitrogen atoms and one sulfur or oxygen atom as ring members,
e.g. 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyrrolyl,
3-pyrrolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl,
3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 3-pyrazolyl,
4-pyrazolyl, 5-pyrazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl,
35 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-imidazolyl,
4-imidazolyl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl,
1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl,
1,2,4-triazol-3-yl, 1,3,4-oxadiazol-2-yl, 1,3,4-thiadiazol-2-yl
and 1,3,4-triazol-2-yl;
40 - 6-membered heteroaryl comprising one to three or one to four
nitrogen atoms: 6-ring heteroaryl groups which, in addition to
carbon atoms, can comprise one to three or one to four nitrogen
atoms as ring members, e.g. 2-pyridinyl, 3-pyridinyl,
4-pyridinyl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrimidinyl,
45 4-pyrimidinyl, 5-pyrimidinyl, 2-pyrazinyl, 1,3,5-triazin-2-yl
and 1,2,4-triazin-3-yl;

14

alkylene: unbranched divalent chains formed from 3 to 5 CH₂ groups, e.g. CH₂, CH₂CH₂, CH₂CH₂CH₂, CH₂CH₂CH₂CH₂ and CH₂CH₂CH₂CH₂CH₂;

5 oxyalkylene: unbranched divalent chains formed from 2 to 4 CH₂ groups in which one valency is bonded to the backbone via an oxygen atom, e.g. OCH₂CH₂, OCH₂CH₂CH₂ and OCH₂CH₂CH₂CH₂;

oxyalkylenoxy: unbranched divalent chains formed from 1 to 3 CH₂ groups in which both valencies is bonded to the backbone via an oxygen atom, e.g. OCH₂O, OCH₂CH₂O and OCH₂CH₂CH₂O.

The (R)- and (S)-isomers and the racemates of compounds of the formula I which have chiral centers are included in the present invention.

The embodiments of the intermediates which are especially preferred with regard to the variables correspond to those of the radicals L¹, L², L³, L⁴, L⁵, R¹, R² and X of the formula I.

20

In view of their intended use of the triazolopyrimidines of the formula I, the following meanings of the substituents, in each case alone or in combination, are especially preferred:

25 Preference is given to compounds I in which R¹ is C₁-C₆-alkyl or C₁-C₈-haloalkyl.


Preference is also given to compounds I in which R¹ is C₂-C₁₀-alkenyl or C₂-C₁₀-alkynyl.

30

Preference is similarly given to compounds I in which R¹ is a saturated or aromatic 5- or 6-membered heterocycle.

Compounds I are particularly preferred in which R¹ is a group B

35



B

in which

40

Y¹ represents hydrogen, fluorine or C₁-C₆-fluoroalkyl,

Y² represents hydrogen or fluorine, or

Y¹ and Y² together form a double bond;

m is 0 or 1; and

45 R³ represents hydrogen or methyl.

15

Preference is furthermore given to compounds I in which R¹ is C₃-C₆-cycloalkyl which can be substituted by C₁-C₄-alkyl.

Preference is particularly given to compounds I in which R² represents hydrogen.

Similarly preferred are compounds I in which R² is methyl or ethyl.

10 If R¹ and/or R² represent halogen-free groups with a center of chirality, the (R)-isomers are preferred. If R¹ and/or R² comprise haloalkyl or haloalkenyl groups with a center of chirality, the (S)-isomers are preferred.

15 Furthermore, particular preference is given to compounds I in which R¹ and R², together with the nitrogen atom to which they are bonded, form a five- or six-membered ring which can be interrupted by an atom from the group consisting of O, N and S and/or can carry one or more substituents from the group
20 consisting of halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl and oxy-C₁-C₃-alkylenoxy or in which an N and a neighboring C atom can be connected via a C₁-C₄-alkylene chain.

Particularly preferred are compounds I in which R¹ and R²,
25 together with the nitrogen atom to which they are bonded, form a 5- or 6-membered ring which optionally can exhibit a double bond and can be substituted as described above.

Particular preference is given in particular to compounds I in
30 which R¹ and R², together with the nitrogen atom to which they are bonded, form a piperidine, morpholine or thiomorpholine ring, in particular a piperidinyl ring, which is optionally substituted by one to three halogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl groups, in particular by 4-methyl.

35 Particularly preferred are furthermore compounds I in which R¹ and R², together with the nitrogen atom to which they are bonded, form a pyrrolidine ring which is optionally substituted by one or two halogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl groups, in particular by
40 2-methyl.

Preferred are compounds of the formula I in which at least one L² and/or L³ group does not represent hydrogen.

45 In addition, compounds of the formula I are preferred in which L¹ represents S(=O)_nA¹, L² represents halogen, L³ and L⁴ represent

16

hydrogen or halogen and L^5 represents hydrogen. They are denoted as compounds I.1.

Similarly preferred are compounds I.1 in which A^1 represents hydrogen or, in particular, methyl.

Preference is given in particular to compounds I.1 in which $n = 0$.

- 10 Preference is given to compounds I.1 in which both L^2 and L^3 represent halogen, in particular fluorine. Furthermore preferred are the compounds I.1 in which L^2 represents fluorine and L^3 represents chlorine or L^2 and L^3 both represent chlorine. L^4 preferably represents hydrogen.

15

An additional preferred embodiment of the compounds of the formula I are those in which L^1 represents cyano or $C(=O)A^2$. They are denoted as compounds I.2.



- 20 Particular preference is furthermore given to compounds I.2 in which L^1 represents C_1 - C_6 -alkoxycarbonyl.

Similarly particularly preferred are compounds I.2 in which L^2 represents halogen and L^3 represents halogen or hydrogen, in


25 particular halogen.

Particular preference is also given to compounds I.2 in which L^4 represents hydrogen and L^5 represents hydrogen or methyl.

- 30 Preference is given in particular to compounds of the formulae IA and IB in which the variables have the meanings given for formula I:



- 40 Preference is furthermore given to compounds IC:



17

Preference is also given, in addition, to compounds I.2 in which L¹ represents C(=O)OCH₃, L² represents fluorine, L³ and L⁵ represent hydrogen and L⁴ represents methyl.

- 5 Particular preference is given to compounds I in which X represents chlorine.

Particular preference is given, in view of their use, to the compounds I compiled in the following tables. The groups

- 10 mentioned in the tables for a substituent additionally represent, considered per se, independently of the combination in which they are mentioned, a particularly preferred embodiment of the substituent in question.

15 Table 1

Compounds of the formula I.1 in which X represents chlorine, L¹ represents methylthio, L² and L³ represent fluorine and L⁴ represents hydrogen and the combination of R¹ and R² for a compound each time corresponds to a row of table A

20

Table 2

Compounds of the formula I.1 in which X represents chlorine, L¹ represents methylsulfinyl, L² and L³ represent fluorine and L⁴ represents hydrogen and the combination of R¹ and R² for a com-

- 25 pound each time corresponds to a row of table A

Table 3

Compounds of the formula I.1 in which X represents chlorine, L¹ represents methylsulfonyl, L² and L³ represent fluorine and L⁴

- 30 represents hydrogen and the combination of R¹ and R² for a compound each time corresponds to a row of table A

Table 4

Compounds of the formula I.1 in which X represents chlorine, L¹

- 35 represents methylthio, L² represents fluorine, L³ represents chlorine and L⁴ represents hydrogen and the combination of R¹ and R² for a compound each time corresponds to a row of table A

Table 5

- 40 Compounds of the formula I.1 in which X represents chlorine, L¹ represents methylsulfinyl, L² represents fluorine, L³ represents chlorine and L⁴ represents hydrogen and the combination of R¹ and R² for a compound each time corresponds to a row of table A

45

18

Table 6

Compounds of the formula I.1 in which X represents chlorine, L¹ represents methylsulfonyl, L² represents fluorine, L³ represents chlorine and L⁴ represents hydrogen and the combination of R¹ and R² for a compound each time corresponds to a row of table A

Table 7

Compounds of the formula I.1 in which X represents chlorine, L¹ represents methylthio, L² and L³ represent chlorine and L⁴ represents hydrogen and the combination of R¹ and R² for a compound each time corresponds to a row of table A

Table 8

Compounds of the formula I.1 in which X represents chlorine, L¹ represents methylsulfinyl, L² and L³ represent chlorine and L⁴ represents hydrogen and the combination of R¹ and R² for a compound each time corresponds to a row of table A


Table 9

Compounds of the formula I.1 in which X represents chlorine, L¹ represents methylsulfonyl, L² and L³ represent chlorine and L⁴ represents hydrogen and the combination of R¹ and R² for a compound each time corresponds to a row of table A

Table 10

Compounds of the formula IA in which X is chlorine, L¹ is C(=O)OCH₃ and L² and L³ are fluorine and the combination of R¹ and R² for a compound each time corresponds to a row of table A

30



IA

35

Table 11

Compounds of the formula IA in which X is chlorine, L¹ is cyano and L² and L³ are fluorine and the combination of R¹ and R² for a compound each time corresponds to a row of table A

40

Table 12

Compounds of the formula IA in which X is chlorine, L¹ is CHO and L² and L³ are fluorine and the combination of R¹ and R² for a compound each time corresponds to a row of table A

45

Table 13

Compounds of the formula IA in which X is chlorine, L¹ is C(=O)CH₃

19

and L^2 and L^3 are fluorine and the combination of R^1 and R^2 for a compound each time corresponds to a row of table A


Table 14

- 5 Compounds of the formula IA in which X is chlorine, L^1 is $C(=O)NHCH_3$ and L^2 and L^3 are [lacuna] and the combination of R^1 and R^2 for a compound each time corresponds to a row of table A

Table 15

- 10 Compounds of the formula IB in which X is chlorine, L^1 is $C(=O)OCH_3$ and L^2 is fluorine and the combination of R^1 and R^2 for a compound each time corresponds to a row of table A

15



IB


20 Table 16

Compounds of the formula IB in which X is chlorine, L^1 is $C(=O)OCH_3$ and L^2 is chlorine and the combination of R^1 and R^2 for a compound each time corresponds to a row of table A

25 Table 17

Compounds of the formula IC in which X is chlorine and L^1 is $C(=O)OCH_3$ and the combination of R^1 and R^2 for a compound each time corresponds to a row of table A

30



IC

35

Table 18

Compounds of the formula IC in which X is chlorine and L^1 is cyano and the combination of R^1 and R^2 for a compound each time corresponds to a row of table A

40

Table 19

Compounds of the formula I in which X is chlorine, L^1 is $C(=O)OCH_3$, and L^2 , L^3 , L^4 and L^5 are fluorine and the combination of R^1 and R^2 for a compound each time corresponds to a row of

45 table A

Table A

No.	R ¹	R ²
5	A-1	CH ₂ CH ₃
	A-2	CH ₂ CH ₃
	A-3	CH ₂ CH ₃
	A-4	CH ₂ CF ₃
	A-5	CH ₂ CF ₃
10	A-6	CH ₂ CF ₃
	A-7	CH ₂ CCl ₃
	A-8	CH ₂ CCl ₃
	A-9	CH ₂ CCl ₃
	A-10	CH ₂ CH ₂ CH ₃
15	A-11	CH ₂ CH ₂ CH ₃
	A-12	CH ₂ CH ₂ CH ₃
	A-13	CH ₂ CH ₂ CH ₃
	A-14	CH(CH ₃) ₂
	A-15	CH(CH ₃) ₂
20	A-16	CH(CH ₃) ₂
	A-17	(±) CH(CH ₃)-CH ₂ CH ₃
	A-18	(±) CH(CH ₃)-CH ₂ CH ₃
	A-19	(±) CH(CH ₃)-CH ₂ CH ₃
	A-20	(S) CH(CH ₃)-CH ₂ CH ₃
25	A-21	(S) CH(CH ₃)-CH ₂ CH ₃
	A-22	(S) CH(CH ₃)-CH ₂ CH ₃
	A-23	(R) CH(CH ₃)-CH ₂ CH ₃
	A-24	(R) CH(CH ₃)-CH ₂ CH ₃
	A-25	(R) CH(CH ₃)-CH ₂ CH ₃
30	A-26	(±) CH(CH ₃)-CH(CH ₃) ₂
	A-27	(±) CH(CH ₃)-CH(CH ₃) ₂
	A-28	(±) CH(CH ₃)-CH(CH ₃) ₂
	A-29	(S) CH(CH ₃)-CH(CH ₃) ₂
	A-30	(S) CH(CH ₃)-CH(CH ₃) ₂
35	A-31	(S) CH(CH ₃)-CH(CH ₃) ₂
	A-32	(R) CH(CH ₃)-CH(CH ₃) ₂
	A-33	(R) CH(CH ₃)-CH(CH ₃) ₂
	A-34	(R) CH(CH ₃)-CH(CH ₃) ₂
	A-35	(±) CH(CH ₃)-C(CH ₃) ₃
40	A-36	(±) CH(CH ₃)-C(CH ₃) ₃
	A-37	(±) CH(CH ₃)-C(CH ₃) ₃
	A-38	(S) CH(CH ₃)-C(CH ₃) ₃

21

No.	R ¹	R ²
A-39	(S) CH(CH ₃)-C(CH ₃) ₃	CH ₃
A-40	(S) CH(CH ₃)-C(CH ₃) ₃	CH ₂ CH ₃
5 A-41	(R) CH(CH ₃)-C(CH ₃) ₃	H
A-42	(R) CH(CH ₃)-C(CH ₃) ₃	CH ₃
A-43	(R) CH(CH ₃)-C(CH ₃) ₃	CH ₂ CH ₃
A-44	(±) CH(CH ₃)-CF ₃	H
10 A-45	(±) CH(CH ₃)-CF ₃	CH ₃
A-46	(±) CH(CH ₃)-CF ₃	CH ₂ CH ₃
A-47	(S) CH(CH ₃)-CF ₃	H
A-48	(S) CH(CH ₃)-CF ₃	CH ₃
A-49	(S) CH(CH ₃)-CF ₃	CH ₂ CH ₃
15 A-50	(R) CH(CH ₃)-CF ₃	H
A-51	(R) CH(CH ₃)-CF ₃	CH ₃
A-52	(R) CH(CH ₃)-CF ₃	CH ₂ CH ₃
A-53	(±) CH(CH ₃)-CCl ₃	H
20 A-54	(±) CH(CH ₃)-CCl ₃	CH ₃
A-55	(±) CH(CH ₃)-CCl ₃	CH ₂ CH ₃
A-56	(S) CH(CH ₃)-CCl ₃	H
A-57	(S) CH(CH ₃)-CCl ₃	CH ₃
25 A-58	(S) CH(CH ₃)-CCl ₃	CH ₂ CH ₃
A-59	(R) CH(CH ₃)-CCl ₃	H
A-60	(R) CH(CH ₃)-CCl ₃	CH ₃
A-61	(R) CH(CH ₃)-CCl ₃	CH ₂ CH ₃
A-62	CH ₂ CF ₂ CF ₃	H
30 A-63	CH ₂ CF ₂ CF ₃	CH ₃
A-64	CH ₂ CF ₂ CF ₃	CH ₂ CH ₃
A-65	CH ₂ (CF ₂) ₂ CF ₃	H
A-66	CH ₂ (CF ₂) ₂ CF ₃	CH ₃
35 A-67	CH ₂ (CF ₂) ₂ CF ₃	CH ₂ CH ₃
A-68	CH ₂ C(CH ₃)=CH ₂	H
A-69	CH ₂ C(CH ₃)=CH ₂	CH ₃
A-70	CH ₂ C(CH ₃)=CH ₂	CH ₂ CH ₃
40 A-71	CH ₂ CH=CH ₂	H
A-72	CH ₂ CH=CH ₂	CH ₃
A-73	CH ₂ CH=CH ₂	CH ₂ CH ₃
A-74	CH(CH ₃)CH=CH ₂	H
45 A-75	CH(CH ₃)CH=CH ₂	CH ₃
A-76	CH(CH ₃)CH=CH ₂	CH ₂ CH ₃
A-77	CH(CH ₃)C(CH ₃)=CH ₂	H

22

No.	R ¹	R ²
A-78	CH(CH ₃)C(CH ₃)=CH ₂	CH ₃
A-79	CH(CH ₃)C(CH ₃)=CH ₂	CH ₂ CH ₃
5 A-80	cyclopentyl	H
A-81	cyclopentyl	CH ₃
A-82	cyclopentyl	CH ₂ CH ₃
A-83	cyclohexyl	H
A-84	cyclohexyl	CH ₃
10 A-85	cyclohexyl	CH ₂ CH ₃
A-86	-(CH ₂) ₂ CH=CHCH ₂ -	
A-87	-(CH ₂) ₂ C(CH ₃)=CHCH ₂ -	
A-88	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -	
15 A-89	-(CH ₂) ₂ CHF(CH ₂) ₂ -	
A-90	-(CH ₂) ₃ CHFCH ₂ -	
A-91	-(CH ₂) ₂ CH(CF ₃)(CH ₂) ₂ -	
A-92	-(CH ₂) ₂ O(CH ₂) ₂ -	
20 A-93	-(CH ₂) ₂ S(CH ₂) ₂ -	
A-94	-(CH ₂) ₅ -	
A-95	-(CH ₂) ₄ -	
A-96	-CH ₂ CH=CHCH ₂ -	
25 A-97	-CH(CH ₃)(CH ₂) ₃ -	
A-98	-CH ₂ CH(CH ₃)(CH ₂) ₂ -	

The compounds I are suitable as fungicides. They are distinguished by an outstanding effectiveness against a broad spectrum of phytopathogenic fungi, especially from the classes of the *Ascomycetes*, *Deuteromycetes*, *Phycomycetes* and *Basidiomycetes*. Some are systemically effective and they can be used in plant protection as foliar and soil fungicides.

They are particularly important in the control of a multitude of fungi on various cultivated plants, such as wheat, rye, barley, oats, rice, maize, grass, bananas, cotton, soya, coffee, sugar cane, vines, fruits and ornamental plants, and vegetables, such as cucumbers, beans, tomatoes, potatoes and cucurbits, and on the seeds of these plants.

They are especially suitable for controlling the following plant diseases:

- *Alternaria* species on fruit and vegetables,
- *Botrytis cinerea* (gray mold) on strawberries, vegetables, ornamental plants and grapevines,
- *Cercospora arachidicola* on peanuts,
- *Erysiphe cichoracearum* and *Sphaerotheca fuliginea* on cucurbits,

23

- *Blumeria graminis* (powdery mildew) on cereals,
- *Fusarium* and *Verticillium* species on various plants,
- *Helminthosporium* species on cereals,
- *Mycosphaerella* species on bananas and peanuts,
- 5 • *Phytophthora infestans* on potatoes and tomatoes,
- *Plasmopara viticola* on grapevines,
- *Podosphaera leucotricha* on apples,
- *Pseudocercospora herpotrichoides* on wheat and barley,
- *Pseudoperonospora* species on hops and cucumbers,
- 10 • *Puccinia* species on cereals,
- *Pyricularia oryzae* on rice,
- *Rhizoctonia* species on cotton, rice and lawns,
- *Septoria nodorum* on wheat,
- *Uncinula necator* on grapevines,
- 15 • *Ustilago* species on cereals and sugar cane, and
- *Venturia* species (scab) on apples and pears.

The compounds I are also suitable for controlling harmful fungi, such as *Paecilomyces variotii*, in the protection of materials

- 20 (e.g. wood, paper, paint dispersions, fibers or fabrics) and in the protection of stored products.

The compounds I are employed by treating the fungi or the plants, seeds, materials or soil to be protected from fungal attack with

25 a fungicidally effective amount of the active compounds. The application can be carried out both before and after the infection of the materials, plants or seeds by the fungi.

- The fungicidal compositions generally comprise between 0.1 and
- 30 95%, preferably between 0.5 and 90%, by weight of active compound.

- When employed in plant protection, the amounts applied are, depending on the kind of effect desired, between 0.01 and 2.0 kg
- 35 of active compound per ha.

In seed treatment, amounts of active compound of 0.001 to 0.1 g, preferably 0.01 to 0.05 g, per kilogram of seed are generally necessary.

40

- When used in the protection of materials or stored products, the amount of active compound applied depends on the kind of application area and on the effect desired. Amounts customarily applied in the protection of materials are, for example, 0.001 g
- 45 to 2 kg, preferably 0.005 g to 1 kg, of active compound per cubic meter of treated material.

24

The compounds I can be converted to the usual formulations, e.g. solutions, emulsions, suspensions, dusts, powders, pastes and granules. The application form depends on the respective use intended; it should in any case guarantee a fine and uniform
5 distribution of the compound according to the invention.

The formulations are prepared in a known way, e.g. by extending the active compound with solvents and/or carriers, if desired using emulsifiers and dispersants, it being possible, when water
10 is the diluent, also to use other organic solvents as auxiliary solvents. Suitable auxiliaries for this purpose are essentially: solvents, such as aromatics (e.g. xylene), chlorinated aromatics (e.g. chlorobenzenes), paraffins (e.g. petroleum fractions), alcohols (e.g. methanol, butanol), ketones (e.g. cyclohexanone),
15 amines (e.g. ethanolamine, dimethylformamide) and water; carriers, such as ground natural minerals (e.g. kaolins, clays, talc, chalk) and ground synthetic ores (e.g. highly dispersed silicic acid, silicates); emulsifiers, such as nonionic and anionic emulsifiers (e.g. polyoxyethylene fatty alcohol ethers,
20 alkylsulfonates and arylsulfonates) and dispersants, such as lignosulfite waste liquors and methylcellulose.

Suitable surfactants are alkali metal, alkaline earth metal and ammonium salts of lignosulfonic acid, naphthalenesulfonic acid,
25 phenolsulfonic acid and dibutylphenylsulfonic acid, alkylarylsulfonates, alkyl sulfates, alkylsulfonates, fatty alcohol sulfates and fatty acids, and alkali metal and alkaline earth metal salts thereof, salts of sulfated fatty alcohol glycol ethers, condensation products of sulfonated naphthalene and
30 naphthalene derivatives with formaldehyde, condensation products of naphthalene or of naphthalenesulfonic acid with phenol and formaldehyde, polyoxyethylene octylphenol ethers, ethoxylated isooctylphenol, octylphenol and nonylphenol, alkylphenol polyglycol ethers, tributylphenyl polyglycol ethers, alkylaryl
35 polyether alcohols, isotridecyl alcohol, fatty alcohol ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers, ethoxylated polyoxypropylene, lauryl alcohol polyglycol ether acetal, sorbitol esters, lignosulfite waste liquors and methylcellulose.

40

Petroleum fractions having medium to high boiling points, such as kerosene or diesel fuel, furthermore coal tar oils, and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, e.g. benzene, toluene, xylene, paraffin,
45 tetrahydronaphthalene, alkylated naphthalenes or derivatives thereof, methanol, ethanol, propanol, butanol, chloroform, carbon tetrachloride, cyclohexanol, cyclohexanone, chlorobenzene or

25

isophorone, or highly polar solvents, e.g. dimethylformamide, dimethyl sulfoxide, N-methylpyrrolidone or water, are suitable for the preparation of directly sprayable solutions, emulsions, pastes or oil dispersions.

5

Powders, combinations for broadcasting and dusts can be prepared by mixing or mutually grinding the active substances with a solid carrier.

- 10 Granules, e.g. coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active compounds to solid carriers. Solid carriers are, e.g., mineral earths, such as silica gels, silicates, talc, kaolin, attaclay, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous
- 15 earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers, such as, e.g., ammonium sulfate, ammonium phosphate, ammonium nitrate or ureas, and plant products, such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders and other solid carriers.

20

The formulations generally comprise between 0.01 and 95% by weight, preferably between 0.1 and 90% by weight, of the active compound. The active compounds are employed therein in a purity of 90% to 100%, preferably 95% to 100% (according to the NMR

25 spectrum).

Examples for formulations are:

- I. 5 parts by weight of a compound according to the invention
- 30 are intimately mixed with 95 parts by weight of finely divided kaolin. In this way, a dust comprising 5% by weight of the active compound is obtained.
- II. 30 parts by weight of a compound according to the invention
- 35 are intimately mixed with a mixture of 92 parts by weight of pulverulent silica gel and 8 parts by weight of liquid paraffin, which had been sprayed onto the surface of this silica gel. In this way, an active compound preparation with good adhesive properties (active compound content 23% by
- 40 weight) is obtained.
- III. 10 parts by weight of a compound according to the invention
- are dissolved in a mixture consisting of 90 parts by weight of xylene, 6 parts by weight of the addition product of 8 to
- 45 10 mol of ethylene oxide with 1 mol of the N-monoethanolamide of oleic acid, 2 parts by weight of the calcium salt of dodecylbenzenesulfonic acid and 2 parts by weight of the

26

addition product of 40 mol of ethylene oxide with 1 mol of castor oil (active compound content 9% by weight).

- IV. 20 parts by weight of a compound according to the invention
5 are dissolved in a mixture consisting of 60 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 5 parts by weight of the addition product of 7 mol of ethylene oxide with 1 mol of isooctylphenol and 5 parts by weight of the
10 addition product of 40 mol of ethylene oxide with 1 mol of castor oil (active compound content 16% by weight).
- V. 80 parts by weight of a compound according to the invention
15 are intimately mixed with 3 parts by weight of the sodium salt of diisobutylnaphthalene- α -sulfonic acid, 10 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 7 parts by weight of pulverulent silica gel and are ground in a hammer mill (active compound content 80% by weight).
- 20 VI. 90 parts by weight of a compound according to the invention are mixed with 10 parts by weight of N-methyl- α -pyrrolidone and a solution is obtained which is suitable for use in the form of very small drops (active compound content 90% by
25 weight).
- VII. 20 parts by weight of a compound according to the invention
30 are dissolved in a mixture consisting of 40 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 20 parts by weight of the addition product of 7 mol of ethylene oxide with 1 mol of isooctylphenol and 10 parts by weight of the
35 addition product of 40 mol of ethylene oxide with 1 mol of castor oil. By running the solution into 100 000 parts by weight of water and finely dispersing it therein, an aqueous dispersion is obtained comprising 0.02% by weight of the active compound.
- VIII. 20 parts by weight of a compound according to the invention
40 are intimately mixed with 3 parts by weight of the sodium salt of diisobutylnaphthalene- α -sulfonic acid, 17 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 60 parts by weight of pulverulent silica gel and are ground in a hammer mill. A spray emulsion comprising 0.1% by weight of the active compound is obtained
45 by fine dispersion of the mixture in 20 000 parts by weight of water.

27

The active compounds can be used as such, in the form of their formulations or of the application forms prepared therefrom, e.g. in the form of directly sprayable solutions, powders, suspensions or dispersions, emulsions, oil dispersions, pastes, dusts, compositions for broadcasting or granules, by spraying, atomizing, dusting, broadcasting or watering. The application forms depend entirely on the intended uses; they should in any case guarantee the finest possible dispersion of the active compounds according to the invention.

10

Aqueous application forms can be prepared from emulsion concentrates, pastes or wettable powders (spray powders, oil dispersions) by addition of water. To prepare emulsions, pastes or oil dispersions, the substances can be homogenized in water, as such or dissolved in an oil or solvent, by means of wetting agents, tackifiers, dispersants or emulsifiers. However, concentrates comprising active substance, wetting agent, tackifier, dispersant or emulsifier and possibly solvent or oil can also be prepared, which concentrates are suitable for dilution with water.

20

The concentrations of active compound in the ready-for-use preparations can be varied within relatively wide ranges. In general, they are between 0.0001 and 10%, preferably between 0.01 and 1%.

25

The active compounds can also be used with great success in the ultra low volume (ULV) process, it being possible to apply formulations with more than 95% by weight of active compound or even the active compound without additives.

30

Oils of various types, herbicides, fungicides, other pesticides and bactericides can be added to the active compounds, if need be also not until immediately before use (tank mix). These agents can be added to the compositions according to the invention in a weight ratio of 1:10 to 10:1.

35

The compositions according to the invention can, in the application form as fungicides, also be present together with other active compounds, which e.g. with herbicides, insecticides, growth regulators, fungicides or also with fertilizers. On mixing the compounds I or the compositions comprising them in the application form as fungicides with other fungicides, in many

45

28

cases an expansion of the fungicidal spectrum of activity is obtained.

The following list of fungicides, with which the compounds according to the invention can be used in conjunction, is intended to illustrate the possible combinations but does not limit them:

- sulfur, dithiocarbamates and their derivatives, such as iron(III) dimethyldithiocarbamate, zinc dimethyldithiocarbamate, zinc ethylenebisdithiocarbamate, manganese ethylenebisdithiocarbamate, manganese zinc ethylenediaminebisdithiocarbamate, tetramethylthiuram disulfide, ammonia complex of zinc (N,N-ethylenebisdithiocarbamate), ammonia complex of zinc (N,N'-propylenebisdithiocarbamate), zinc (N,N'-propylenebisdithiocarbamate) or N,N'-polypropylenebis(thiocarbamoyl)disulfide;
- nitro derivatives, such as dinitro(1-methylheptyl)phenyl crotonate, 2-sec-butyl-4,6-dinitrophenyl 3,3-dimethylacrylate, 2-sec-butyl-4,6-dinitrophenyl isopropyl carbonate or diisopropyl 5-nitroisophthalate;
- heterocyclic substances, such as 2-heptadecyl-2-imidazoline acetate, 2-chloro-N-(4'-chlorobiphenyl-2-yl)nicotinamide, 2,4-dichloro-6-(o-chloroanilino)-s-triazine, O,O-diethyl phthalimidophosphonothioate, 5-amino-1-[bis(dimethylamino)phosphinyl]-3-phenyl-1,2,4-triazole, 2,3-dicyano-1,4-dithioanthraquinone, 2-thio-1,3-dithiolo[4,5-b]quinoxaline, methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate, 2-(methoxycarbonylamino)benzimidazole, 2-(2-furyl)benzimidazole, 2-(4-thiazolyl)benzimidazole, N-(1,1,2,2-tetrachloroethylthio)tetrahydrophthalimide, N-(trichloromethylthio)tetrahydrophthalimide or N-(trichloromethylthio)phthalimide,
- N-dichlorofluoromethylthio-N',N'-dimethyl-N-phenylsulfamide, 5-ethoxy-3-trichloromethyl-1,2,3-thiadiazole, 2-thiocyanatomethylthiobenzothiazole, 1,4-dichloro-2,5-dimethoxybenzene, 4-(2-chlorophenylhydrazono)-3-methyl-5-isoxazolone, 2-thiopyridine 1-oxide, 8-hydroxyquinoline or its copper salt, 2,3-dihydro-5-carboxanilido-6-methyl-1,4-oxathiin, 2,3-dihydro-5-carboxanilido-6-methyl-1,4-oxathiin 4,4-dioxide, 2-methyl-5,6-dihydro-4H-pyran-3-carboxanilide, 2-methylfuran-3-carboxanilide, 2,5-dimethylfuran-3-carboxanilide, 2,4,5-trimethylfuran-3-carboxanilide, N-cyclohexyl-2,5-dimethylfuran-3-carboxamide,

29

- N-cyclohexyl-N-methoxy-2,5-dimethylfuran-3-carboxamide,
2-methylbenzanilide, 2-iodobenzanilide, N-formyl-N-morpholine
2,2,2-trichloroethyl acetal, piperazin-1,4-diylbis-1-
(2,2,2-trichloroethyl)formamide,
- 5 1-(3,4-dichloroanilino)-1-formylamino-2,2,2-trichloroethane,
2,6-dimethyl-N-tridecylmorpholine or its salts,
2,6-dimethyl-N-cyclododecylmorpholine or its salts,
N-[3-(p-(tert-butyl)phenyl)-2-methylpropyl]-cis-2,6-dimethyl-
morpholine, N-[3-(p-(tert-butyl)phenyl)-2-methyl
- 10 propyl]piperidine, 1-[2-(2,4-dichlorophenyl)-4-ethyl-
1,3-dioxolan-2-ylethyl]-1H-1,2,4-triazole,
1-[2-(2,4-dichlorophenyl)-4-(n-propyl)-1,3-dioxolan-2-yl-
ethyl]-1H-1,2,4-triazole,
N-(n-propyl)-N-(2,4,6-trichlorophenoxyethyl)-N'-imidazolylurea,
- 15 1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-
butanone,
1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-
butanol,
(2RS,3RS)-1-[3-(2-chlorophenyl)-2-(4-fluorophenyl)oxiran-2-
ylmethyl]-1H-1,2,4-triazole,
- 20 α -(2-chlorophenyl)- α -(4-chlorophenyl)-5-pyrimidinemethanol,
5-butyl-2-dimethylamino-4-hydroxy-6-methylpyrimidine,
bis(p-chlorophenyl)-3-pyridinemethanol,
1,2-bis(3-ethoxycarbonyl-2-thioureido)benzene or
- 25 1,2-bis(3-methoxycarbonyl-2-thioureido)benzene,
• strobilurins, such as methyl
E-methoxyimino[α -(o-tolyloxy)-o-tolyl]acetate, methyl
E-2-[2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl]-3-
methoxyacrylate, methyl-E-methoxyimino-[α -(2-phenoxyphenyl)]-
30 acetamide, methyl-E-methoxyimino[α -(2,5-dimethylphenoxy)-
o-tolyl]acetamide, methyl E-2-[2-[2-trifluoromethylpyrid-6-
yl]oxymethyl]phenyl]-3-methoxyacrylate, methyl (E,E)-methoxy
imino{2-[1-(3-trifluoromethylphenyl)ethylideneaminooxymethyl]
phenyl}acetate or methyl N-(2-[1-(4-chlorophenyl)-1H-pyrazol-
35 3-yl]oxymethyl)phenyl)-N-methoxycarbamate,
• anilinopyrimidines, such as
N-(4,6-dimethylpyrimidin-2-yl)aniline,
N-[4-methyl-6-(1-propynyl)pyrimidin-2-yl]aniline or
N-[4-methyl-6-cyclopropylpyrimidin-2-yl]aniline,
- 40 • phenylpyrroles, such as
4-(2,2-difluoro-1,3-benzodioxol-4-yl)pyrrole-3-carbonitrile,
• cinnamamides, such as
3-(4-chlorophenyl)-3-(3,4-dimethoxyphenyl)acryloylmorpholine
or 3-(4-fluorophenyl)-3-(3,4-dimethoxyphenyl)acryloylmorpholine,
- 45 • and various fungicides, such as dodecylguanidine acetate,
1-(3-bromo-6-methoxy-2-methylphenyl)-1-(2,3,4-trimethoxy-6-

30

- methylphenyl)methanone, 3-[3-(3,5-dimethyl-2-oxycyclohexyl)-2-hydroxyethyl]glutarimide,
hexachlorobenzene,
methyl N-(2,6-dimethylphenyl)-N-(2-furoyl)-DL-alaninate,
5 N-(2,6-dimethylphenyl)-N-(2'-methoxyacetyl)-DL-alanine methyl ester, N-(2,6-dimethylphenyl)-N-chloroacetyl-D,L-2-aminobutyrolactone, N-(2,6-dimethylphenyl)-N-(phenylacetyl)-DL-alanine methyl ester, 5-methyl-5-vinyl-3-(3,5-dichlorophenyl)-2,4-dioxo-1,3-
10 oxazolidine, 3-(3,5-dichlorophenyl)-5-methyl-5-methoxymethyl-1,3-oxazolidine-2,4-dione, 3-(3,5-dichlorophenyl)-1-isopropylcarbamoylhydantoin,
N-(3,5-dichlorophenyl)-1,2-dimethylcyclopropane-1,2-
15 dicarboximide,
2-cyano-[N-(ethylaminocarbonyl)-2-methoxyimino]acetamide,
1-[2-(2,4-dichlorophenyl)pentyl]-1H-1,2,4-triazole,
2,4-difluoro- α -(1H-1,2,4-triazolyl-1-methyl)benzhydryl alcohol,
N-(3-chloro-2,6-dinitro-4-trifluoromethylphenyl)-5-trifluoro-
20 methyl-3-chloro-2-aminopyridine, 1-((bis(4-fluorophenyl)methylsilyl)methyl)-1H-1,2,4-triazole,
5-chloro-2-cyano-4-(p-tolyl)imidazole-1-sulfonic acid dimethylamide
25 or 3,5-dichloro-N-(3-chloro-1-ethyl-1-methyl-2-oxopropyl)-4-methylbenzamide.

Synthesis examples

- 30 The procedures described in the following synthesis examples were used to prepare further compounds I by appropriate modification of the starting compounds. The compounds thus obtained are listed in the following table, together with physical data.

35 Example 1: Preparation of diethyl 4-cyanophenylmalonate

- Diethyl malonate (0.49 mol) was added at approximately 60°C over 2 hours to a suspension of sodium hydride (0.51 mol) in 140 ml of 1,4-dioxane. After stirring for a further 10 min, 0.05 mol of
40 CuBr were added. After 15 min, 0.25 mol of 4-cyanobromobenzene in 10 ml of dioxane were added. The reaction mixture was maintained at 100°C for approximately 14 hours and then, at approximately 15°C, 35 ml of 12N hydrochloric acid were slowly added. The precipitate was filtered off and the filtrate was extracted with
45 diethyl ether. After phase separation, the organic phase was

31

dried and then freed from the solvent. 32 g of the title compound remained.

Example 2: Preparation of

5 5,7-dihydroxy-6-(4-cyanophenyl)-[1,2,4]triazolo[1,5-a]pyrimidine

A mixture of 14 g of 3-amino-1,2,4-triazole, 0.17 mol of the ester from example 1 and 50 ml of tributylamine (50 ml) was stirred at 180°C for approximately 6 hours. A solution of 21 g of NaOH in 200 ml of water was added at approximately 70° and the mixture was stirred for a further 30 min. The organic phase was separated and the aqueous phase was extracted with diethyl ether. The product precipitated from the aqueous phase after acidification with concentrated hydrochloric acid. 28 g of the title compound were obtained by filtration.

Example 3: Preparation of

5,7-dichloro-6-(4-cyanophenyl)-[1,2,4]triazolo[1,5-a]pyrimidine

A mixture of 25 g of the triazolopyrimidine from example 2 and 50 ml of POCl₃ was refluxed for 8 hours, POCl₃ being distilled off. The residue was added to a CH₂Cl₂-water mixture and the organic phase was separated, washed, dried and then freed from the solvent. 23 g of the title compound were obtained.

25

Example 4: Preparation of

5-chloro-6-(4-cyanophenyl)-7-isopropylamino-[1,2,4]triazolo[1,5-a]pyrimidine [I-1]

A solution of 1.5 mmol of isopropylamine and 1.5 mmol of triethylamine in 10 ml of dichloromethane was added with stirring to a solution of 1.5 mmol of the product from ex. 3 in 20 ml of dichloromethane. The reaction mixture was stirred at 20 to 25°C for approximately 16 hours and then washed with dilute hydrochloric acid. The organic phase was separated, dried and freed from the solvent. After chromatography on silica gel, 330 mg of the title compound with a melting point of 190°C were obtained.

40 Example 5: Preparation of

5,7-dihydroxy-6-(2,6-difluoro-4-thiomethylphenyl)-[1,2,4]triazolo[1,5-a]pyrimidine

A mixture of 3-amino-1,2,4-triazole (14 g), diethyl (2,6-difluoro-4-(thiomethyl)phenyl)malonate (0.17 mol) and tributylamine (50 ml) was heated at 180°C for approximately 6 hours. After cooling the reaction mixture to 70°C, the solution

45

32

was treated with 21 g of NaOH in 200 ml of water and stirred for a further 30 min. After separating the organic phase and extracting the aqueous phase with diethyl ether, the title compound was precipitated from the aqueous phase by acidification with concentrated HCl solution. 37 g were isolated.

Example 6: Preparation of
5,7-dichloro-6-(2,6-difluoro-4-(thiomethyl)phenyl)-[1,2,4]
triazolo[1,5-a]pyrimidine

10

A mixture of
5,7-dihydroxy-6-(2,6-difluoro-4-(thiomethyl)phenyl)-[1,2,4]
triazolo[1,5-a]pyrimidine (30 g, Ex. 5) and 50 ml of POCl₃ was
refluxed for approximately eight hours; some POCl₃ was distilled
15 off in the process. The residue was added to a CH₂Cl₂/water
mixture. The organic phase was separated, dried and freed from
the solvent. 21 g of the title compound with a melting point of
138°C remained behind.

20 Example 7: Preparation of
5-chloro-6-(2,6-difluoro-4-(thiomethyl)phenyl)-7-(1,1,1-
trifluoroprop-2-yl)amino-[1,2,4]triazolo[1,5-a]pyrimidine [I-6]

A solution of 6 mmol of
25 5,7-dichloro-6-(2,6-difluoro-4-(thiomethyl)phenyl)-[1,2,4]
triazolo[1,5-a]pyrimidine (Ex. 6) in 20 ml of dichloromethane was
treated, with stirring, with a solution of 6 mmol of
1,1,1-trifluoro-2-aminopropane and 6 mmol of triethylamine in
40 ml of dichloromethane. The solution was stirred at 20-25°C for
30 approximately 16 hours and then was washed with dilute HCl
solution. The organic phase was separated, dried and freed from
the solvent. After chromatography on silica gel, 1.2 g of the
title compound with a melting point of 174°C were obtained from
the residue.

35

Example 8: Preparation of
5-chloro-6-(2,6-difluoro-4-(methylsulfonyl)phenyl)-7-(1,1,1-
trifluoroprop-2-yl)amino-[1,2,4]triazolo[1,5-a]pyrimidine (8a)
and 5-chloro-6-(2,6-difluoro-4-(methylsulfinyl)phenyl)-7-
40 (1,1,1-trifluoroprop-2-yl)amino-[1,2,4]triazolo[1,5-a]pyrimidine
(8b)

A solution of 3 mmol of
5-chloro-6-(2,6-difluoro-4-(thiomethyl)phenyl)-7-(1,1,1-
45 trifluoroprop-2-yl)amino-[1,2,4]triazolo[1,5-a]pyrimidine (Ex. 7)
in 20 ml of dichloromethane was treated with 0.13 g of ammonium
molybdate and 0.22 ml of 98% formic acid. 9 mmol of H₂O₂ were

33

added and then the reaction mixture was stirred at 20-25°C for 24 hours. The reaction mixture was added to water and the organic phase was separated, washed with 10% NaHSO₃ solution, dried and freed from the solvent. After chromatography on silica gel, 0.28 g of the sulfone (8a) with a melting point of 211°C and 0.39 g of the sulfoxide (8b) with a melting point of 264°C were obtained.

Example 9: Preparation of 5-cyano-6-(2,6-difluoro-4-cyanophenyl)-7-(4-methylpiperidin-1-yl)-[1,2,4]triazolo[1,5-a]pyrimidine

A mixture of 0.1 mol of 5-chloro-6-(2,6-difluoro-4-cyanophenyl)-7-(4-methylpiperidin-1-yl)-[1,2,4]triazolo[1,5-a]pyrimidine (No. I-5) and 0.25 mol of tetraethylammonium cyanide in 750 ml of dimethylformamide (DMF) was stirred at 20-25°C for 16 hours. After addition of water and methyl tert-butyl ether (MTBE), the organic phase was separated, washed with water, dried and freed from the solvent. After chromatography on silica gel, 6.33 g of the title compound were obtained from the residue.

¹H NMR: 8.55 (s), 7.45 (d), 3.80 (d), 2.95 (t), 1.70 (m), 1.65 (m), 1.40 (m), 0.98 (d).

Example 10: Preparation of 5-methoxy-6-(2,6-difluoro-4-cyanophenyl)-7-(4-methylpiperidin-1-yl)-[1,2,4]triazolo[1,5-a]pyrimidine

A solution of 65 mmol of compound I-5 in 400 ml of anhydrous methanol was, after addition of 71.5 mmol of a 30% sodium methoxide solution at 20-25°C, stirred at this temperature for approximately 16 hours. Methanol was distilled off and the residue was taken up in dichloromethane, then washed with water and, after drying, freed from the solvent. After chromatography on silica gel, 3.68 g of the title compound were obtained from the residue.

¹H NMR: 8.75 (s), 7.35 (d), 3.95 (s), 3.65 (d), 2.70 (t), 1.65 (m), 1.55 (m), 1.45 (m), 0.95 (d).

Example 11: Preparation of 5-methyl-6-(2,6-difluoro-4-cyanophenyl)-7-(4-methylpiperidin-1-yl)-[1,2,4]triazolo[1,5-a]pyrimidine

A mixture of 20 ml of diethyl malonate and 0.27 g (5.65 mmol) of NaH (50% dispersion in mineral oil) in 50 ml of acetonitrile was stirred at 20-25°C for approximately 2 hours. 4.71 mmol of compound I-5 were added to this mixture and then the mixture was heated to approximately 60°C and stirred at this temperature for

34

20 hours. After addition of 50 ml of aqueous NH_4Cl solution, the acidification was carried out with dilute HCl . After extracting with MTBE, the combined organic phases were dried and evaporated. The residue was purified by chromatography on silica gel. The
5 pure product was taken up in concentrated HCl and held at 80°C for 24 hours. After cooling the reaction mixture, the pH was adjusted to 5 by addition of aqueous NaOH solution and then the mixture was extracted with MTBE. The combined organic phases were, after drying, freed from the solvent. After chromatography on silica
10 gel, 0.78 g of the title compound was obtained from the residue.

^1H NMR: 8.75 (s), 7.35 (d), 3.65 (d), 2.70 (t), 2.43 (s), 1.65 (m), 1.55 (m), 1.45 (m), 0.95 (d).

15 Example 12: Preparation of
5-chloro-6-(2,6-difluoro-4-carboxaldehydephenyl)-7-(4-methylpiperidin-1-yl)-[1,2,4]triazolo[1,5-a]pyrimidine

A solution of 1.5 mmol of compound I-5 in 20 ml of
20 dichloromethane was treated at 0°C with a 1M solution of 1.65 mmol of diisobutylaluminum hydride (DIBAH) in dichloromethane and was stirred at $20-25^\circ\text{C}$ for 2 hours. This mixture was treated with saturated NH_4Cl solution and 10% HCl solution and the organic phase was separated and washed with water. After drying, the
25 solvent was removed and, after chromatography on silica gel, 0.36 g of the title compound was obtained from the residue.

^1H NMR: 10.05 (s), 8.40 (s), 7.60 (d), 3.70 (d), 2.85 (t), 1.65 (m), 1.55 (m), 1.40 (m), 0.95 (d).


30

Example 13: Preparation of 5-chloro-6-(2,6-difluoro-4-acetylphenyl)-7-(4-methylpiperidin-1-yl)-[1,2,4]triazolo[1,5-a]pyrimidine

A solution of 1.5 mmol of compound I-5 in 20 ml of
35 tetrahydrofuran (THF) was treated, at $20-25^\circ\text{C}$, with 1.65 mmol of CuBr and a 3M solution of 1.65 mmol of methylmagnesium chloride in THF and the mixture was stirred for approximately 30 min. This mixture was treated with saturated NH_4Cl solution and 10% HCl solution and the organic phase was separated and washed with
40 water. After drying, the solvent was removed and, after chromatography on silica gel, 0.22 g of the title compound was obtained from the residue.

^1H NMR: 8.40 (s), 7.65 (d), 3.70 (d), 2.80 (t), 2.70 (s),
45 1.70 (m), 1.55 (m), 1.40 (m), 0.98 (d).

Table I



I

No.	R ¹	R ²	L ¹	L ²	L ³	L ⁴	L ⁵	X	Phys. data (M.p. [°C])
I-1	CH(CH ₃) ₂	H	CN	H	H	H	H	Cl	190
I-2	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -		COOCH ₃	F	F	H	H	Cl	8.7(s), 7.8(d), 3.9(s), 3.7(d), 2.8(t), 1.6(m), 1.2(m), 0.9(d)
I-3	cyclopentyl	H	CN	H	H	H	H	Cl	207
I-4	CH(CH ₃)C(CH ₃) ₃	H	CN	H	H	H	H	Cl	130
I-5	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -		CN	F	F	H	H	Cl	8.4(s), 7.4(d), 3.7(d), 2.8(t), 1.7(m), 1.4(m), 0.9(d)
I-6	CH(CH ₃)-CF ₃	H	SCH ₃	F	F	H	H	Cl	174
I-7	CH(CH ₃)-CF ₃	H	SCH ₃	F	H	F	H	Cl	164
I-8	CH ₂ CF ₃	H	SCH ₃	F	H	F	H	Cl	161

In the case of chiral R¹ groups, because of the hindered rotation of the phenyl group, two diastereoisomers A) and B) may exist, which may differ in their physical properties.

Examples for the action against harmful fungi

The fungicidal action of the compounds of the general formula I
5 can be demonstrated from the following tests:

The active compounds were prepared, separately or together, as a
10% emulsion in a mixture of 70% by weight of cyclohexanone, 20%
by weight of Nekanil® LN (Lutensol® AP6, wetting agent with an
10 emulsifying and dispersing action based on ethoxylated
alkylphenols) and 10% by weight of Wettol® EM (nonionic
emulsifier based on ethoxylated castor oil) and were
appropriately diluted with water to the desired concentration.

15 Use example 1 - Activity against early blight of tomato caused by *Alternaria solani*

Leaves of pot plants of the variety "Große Fleischtomate St.
Pierre" were sprayed to runoff point with an aqueous suspension
20 prepared from a stock solution consisting of 10% of active
compound, 85% of cyclohexanone and 5% of emulsifier. On the
following day, the leaves were infected with an aqueous
suspension of zoospores of *Alternaria solani* in 2% Biomalz
solution with a concentration of 0.17×10^6 spores/ml. The plants
25 were subsequently placed in a chamber saturated with water vapor
at temperatures between 20 and 22°C. After 5 days, early blight in
the untreated but infected control plants had so extensively
developed that the infection could be visually determined in %.

30 In this test, the plants treated with 63 ppm of the active
compound I-2 of table I showed no infection, while the untreated
plants were 90% infected.

35 Use example 2: Activity against net blotch of barley caused by *Pyrenophora teres*

Leaves of pot-grown barley seedlings of the variety "Igri" were
sprayed to runoff point with an aqueous preparation of active
compound prepared from a stock solution consisting of 10% of
40 active compound, 85% of cyclohexanone and 5% of emulsifier. 24
hours after the spray coating had dried on, the leaves were
inoculated with an aqueous suspension of spores of *Pyrenophora*
[syn. *Drechslera*] *teres*, the causative agent of net blotch. The
test plants were subsequently placed in a greenhouse at
45 temperatures of between 20 and 24°C and a relative atmospheric
humidity of 95 to 100%. After 6 days, the extent of development

37

of the disease was determined visually in % of infection of the total leaf area.

In this test, the plants treated with 16 ppm of the active compound I-2 of table I showed 1% infection, while the untreated plants were 85% infected.

Use example 3 - Activity against early blight of tomato caused by *Alternaria solani*

10

Pot plants of the variety "Pixie II" in the 2-4-leaf stage were sprayed to runoff point with an aqueous suspension which comprised the active compound in the concentration given below and which was prepared from a stock solution of 5% of active compound, 94% of cyclohexanone and 1% of emulsifier (Tween 20™). After the leaves had dried (3 to 5 hours), they were infected with an aqueous suspension of zoospores of *Alternaria solani* comprising 0.15×10^3 spores/ml. The plants were subsequently placed in a chamber saturated with water vapor at temperatures between 22 and 24°C for 36 hours and then in a greenhouse at a relative humidity of 95% at temperatures between 21 and 23°C for 2 to 3 days. The extent of the development of infection on the leaf undersides was then determined visually.

25 In this test, the plants treated with 200 ppm of the active compounds No. I-6, I-7 and I-8 showed no infection in excess of 7%, while the untreated plants were 100% infected.

Use example 4 - Activity against *Rhizoctonia solani* on rice

30

Pot plants of the variety "M-202" in the 2-leaf stage were sprayed to runoff point with an aqueous preparation of active compound prepared with a stock solution of 5% of active compound, 94% of cyclohexanone and 1% of emulsifier (Tween 20™). After the leaves had dried (3 to 5 hours), they were inoculated, 4 ml of an aqueous mycelium suspension of *Rhizoctonia solani* being pipetted onto the soil surface of each and every pot. The plants were subsequently placed in a chamber saturated with water vapor at temperatures between 22 and 24°C for 36 hours and then in a greenhouse at a relative humidity of 95% at temperatures between 21 and 23°C for 2 to 3 days. The extent of the development of infection on the leaf undersides was then determined visually. The test plants were subsequently placed in climatic chambers at 18-28°C and high atmospheric humidity for 4-5 days. The extent of the development of infection on the leaves was then determined visually.

38

In this test, the plants treated with 200 ppm of the active compounds No. I-6, I-7 and I-8 did not show more than 7% infection, while the untreated plants were 100% infected.

5

10

15

20

25

30

35

40


45

39

We claim:

1. A triazolopyrimidine of the formula I

5



I

10

in which substituents have the following meanings:

L¹ is cyano, S(=O)_nA¹ or C(=O)A², wherein

15

A¹ is hydrogen, hydroxyl, C₁-C₈-alkyl, C₁-C₈-alkylamino or di(C₁-C₈-alkyl)amino;

A² is C₁-C₈-alkoxy, C₁-C₆-haloalkoxy or one of the groups mentioned under A¹;

20

n is 0, 1 or 2;

L², L³ are hydrogen or halogen, wherein, when L¹ is S(=O)A¹, L² is halogen;

25

L⁴, L⁵ are hydrogen, halogen or C₁-C₄-alkyl;

X is halogen, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₂-haloalkoxy;

30

R¹ is C₁-C₈-alkyl, C₁-C₈-haloalkyl, C₃-C₆-cycloalkyl, C₃-C₆-halocycloalkyl, C₂-C₈-alkenyl, C₄-C₁₀-alkadienyl, C₂-C₈-haloalkenyl, C₃-C₆-cycloalkenyl, C₂-C₈-alkynyl, C₂-C₈-haloalkynyl or C₃-C₆-cycloalkynyl, phenyl, naphthyl, or a five- to ten-membered saturated, partially unsaturated or aromatic heterocycle, comprising one to four heteroatoms from the group consisting of O, N and S;

35

40

R² is hydrogen or one of the groups mentioned under R¹,

45

R¹ and R² can also, together with the nitrogen atom to which they are bonded, form a five- or six-membered ring which can be interrupted by an atom from the group consisting of O, N and S and/or can carry one or more substituents from the group consisting of halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl and oxy-C₁-C₃-alkylenoxy or

40

in which an N and a neighboring C atom can be connected via a C₁-C₄-alkylene chain;

wherein R¹ and/or R² can be substituted by one to four identical or different groups R^a:

5

R^a is halogen, cyano, nitro, hydroxyl, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkylcarbonyl, C₃-C₆-cycloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₁-C₆-alkoxycarbonyl, C₁-C₆-alkylthio, C₁-C₆-alkylamino, di(C₁-C₆-alkyl)amino, C₂-C₆-alkenyl, C₂-C₆-alkenyloxy, C₃-C₆-alkynyloxy, C₃-C₆-cycloalkyl, phenyl, naphthyl or a five- to ten-membered saturated, partially unsaturated or aromatic heterocycle, comprising one to four heteroatoms from the group consisting of O, N and S,

10

15

20

wherein these aliphatic, alicyclic or aromatic groups, for their part, can be partially or completely halogenated or can carry one to three groups R^b:

25

30

35

R^b is halogen, cyano, nitro, hydroxyl, mercapto, amino, carboxyl, aminocarbonyl, aminothiocarbonyl, alkyl, haloalkyl, alkenyl, alkenyloxy, alkynyloxy, alkoxy, haloalkoxy, alkylthio, alkylamino, dialkylamino, formyl, alkylcarbonyl, alkylsulfonyl, alkylsulfoxyl, alkoxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, dialkylaminocarbonyl, dialkylaminothiocarbonyl or dialkylaminothiocarbonyl, wherein the alkyl groups in these radicals comprise 1 to 6 carbon atoms and the alkenyl or alkynyl groups mentioned in these radicals comprise 2 to 8 carbon atoms;

40

and/or one to three of the following radicals:

45

cycloalkyl, cycloalkoxy, heterocyclyl or heterocyclyloxy, wherein the cyclic systems comprise 3 to 10 ring members; aryl, aryloxy, arylthio, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, hetaryl, hetaryloxy or hetarylthio, wherein the aryl radicals preferably comprise 6 to 10 ring members and the hetaryl radicals

41

comprise 5 or 6 ring members, wherein the cyclic systems can be partially or completely halogenated or can be substituted by alkyl or haloalkyl groups,

5

A¹ is hydrogen, hydroxyl, C₁-C₈-alkyl, C₁-C₈-alkylamino or di(C₂-C₈-alkyl)amino; and

n is 0, 1 or 2.

10

2. A compound of the formula I as claimed in claim 1, in which

L¹ represents S(=O)_nA¹;

15

L² represents halogen;

L³, L⁴ represent hydrogen or halogen; and

L⁵ represents hydrogen.

20

3. A compound of the formula I as claimed in claim 1, in which

L¹ represents cyano or C(=O)A².

25

4. A compound of the formula I as claimed in any of claims 1 to 3, in which X represents halogen.

5. A compound of the formula I as claimed in any of claims 1 to 4, in which R¹ and R² have the following meanings:

30

R¹ is C₁-C₆-alkyl, C₁-C₈-haloalkyl, C₃-C₆-cycloalkyl, C₃-C₆-halocycloalkyl, C₂-C₈-alkenyl, C₂-C₈-haloalkenyl, C₂-C₈-alkynyl; and

35

R² is hydrogen or C₁-C₄-alkyl; or


R¹ and R² can also, together with the nitrogen atom to which they are bonded, form a saturated or unsaturated 5- or 6-membered ring which can carry one or two substituents from the group consisting of halogen, C₁-C₆-alkyl and C₁-C₆-haloalkyl.

40

45


42

6. A process for the preparation of the compounds of the formula I as claimed in claim 1 in which X is halogen, cyano, C₁-C₄-alkoxy or C₁-C₂-haloalkoxy, by reaction of 5-aminotriazole of the formula II




II

with phenylmalonates of the formula III,




III

to give dihydroxytriazolopyrimidines of the formula IV




IV

and halogenation to give the dihalogen compounds of the formula V,



V

in which Y is halogen, in particular chlorine or bromine, reaction with amines of the formula VI,



VI

in which R¹ and R² have the meanings given in claim 1, to give 5-halo-7-aminotriazolopyrimidines of the formula I in which X is halogen, and, for the preparation of compounds of the formula I in which X represents cyano, C₁-C₄-alkoxy or C₁-C₂-haloalkoxy, reaction with compounds of the formula VII,




VII

which, according to the meaning of the group X' to be introduced, represents an inorganic cyanide, alkoxide or haloalkoxide and in which M is an ammonium,


43

tetraalkylammonium, alkali metal or alkaline earth metal cation.


7. A process for the preparation of the compounds of the formula I as claimed in claim 1 in which X is C₁-C₄-alkyl, by reaction of 5-aminotriazole of the formula II as claimed in claim 6 with dicarbonyl compounds of the formula IIIa,



- 15 in which R and X¹ are C₁-C₄-alkyl, to give hydroxytriazolopyrimidines of the formula IVa



halogenation to give compounds of the formula V,



- 30 in which Y is halogen, especially chlorine or bromine, and reaction with amines of the formula VI as claimed in claim 6 to give triazolopyrimidines of the formula I in which X is C₁-C₄-alkyl.

- 35 8. A compound of the formulae IIIa, IV, IVa, V and Va as claimed in claim 6 or 7.

- 40 9. A composition suitable for the control of harmful fungi, comprising a solid or liquid carrier and a compound of the formula I as claimed in claim 1.

- 45 10. A method for the control of harmful fungi, which comprises treating the fungi or the materials, plants, ground or seeds to be protected from fungal attack with an effective amount of a compound of the formula I as claimed in claim 1.